

# СПИСОК ЛИТЕРАТУРЫ

*Адрианов О.С., Выходцева Н.И., Фокин В.Ф. и др.* Метод локального воздействия фокусированным ультразвуком на глубоко расположенные структуры мозга необездвиженного ненаркотизированного животного. Бюллетень эксперимент. биол. и мед. 1984а; 98(7): 115–117.

*Адрианов О.С., Выходцева Н.И., Фокин В.Ф. и др.* Обратимое функциональное выключение зрительного тракта при действии фокусированным ультразвуком. Бюллетень эксперимент. биол. и мед. 1984б; 97(6): 760–762.

*Алтухов Н.В.* Энцефалометрические исследования мозга в связи с полом, возрастом и черепным указателем. М.: Университетская типография; 1891. 56 с.

*Ахмадеева Г.Н., Галимова Р.М., Набиуллина Д.И. и др.* Лечение акинетико-ригидной формы болезни Паркинсона, осложненной развитием флуктуаций и дискинезий, методом деструкции с помощью фокусированного ультразвука. Нервные болезни. 2022; 4: 26–31.

*Бриль Е.В.* Ранняя нейростимуляция подкорковых структур головного мозга при болезни Паркинсона. Споры продолжаются. Бюллетень Национального общества по изучению болезни Паркинсона и расстройств движений. 2020; 3: 13–15.

*Бриль Е.В., Белова Е.М., Седов А.С. и др.* Современные представления о механизмах нейростимуляции при болезни Паркинсона. Анн. клин. и эксперимент. неврол. 2022; 16(2): 89–99.

*Васин Н.Я., Меджидов М.Р., Шабалов В.А.* Стереотаксическая деструкция интерстициального ядра Кахала при спастической кривошее. Журн. «Вопросы нейрохирургии» им. Н.Н. Бурденко. 1985; 4: 3–7.

Высший титул. Воспоминания о докторе Э. Канделе. М.: Эллис Лак; 1994. 155 с.

*Галимова Р.М., Иллариошкин С.Н., Бузаев И.В., Качемаева О.В.* Терапия двигательных нарушений методом фокусированного ультразвука под контролем магнитно-резонансной томографии. Рекомендации для врачей-неврологов по отбору пациентов. Бюллетень Национального общества по изучению болезни Паркинсона и расстройств движений. 2020; 1: 9–15.

*Галимова Р.М., Набиуллина Д.И., Иллариошкин С.Н. и др.* Первый в России опыт лечения пациентов с эссенциальным тремором методом фокусированного ультразвука под контролем МРТ. Анн. клин. и эксперимент. неврол. 2022; 16(2): 5–14.

*Галимова Р.М., Сидорова Ю.А., Набиуллина Д.И. и др.* Фокусированный ультразвук под контролем МРТ в лечении цервикальной дистонии. Анн. клин. и эксперимент. неврол. 2023; 17(4): 28–34.

*Гурбатов С.Н., Клемина А.В.* Влияние радиационной силы на процесс оседания эритроцитов. Вестник Нижегородского университета им. Н.И. Лобачевского. 2011; 5-3: 136–141.

*Гуца А.О., Тюрников В.М., Кащеев А.А.* Современные возможности хирургической нейромодуляции. Анн. клин. и эксперимент. неврол. 2018; 12(спец. вып.): 32–37.

*Давыдов О.С., Яхно Н.Н., Кукушкин М.Л. и др.* Невропатическая боль: клинические рекомендации по диагностике и лечению Российского общества по изучению боли. Российский журн. боли. 2018; 4: 5–41.

*Зернов Д.Н.* Энцефалометр: прибор для определения положения частей мозга у живого человека. М.: Печатня С.П. Яковлева; 1892.

*Иллариошкин С.Н.* Конформационные болезни мозга. М.: Янус-К; 2003. 248 с.

*Иллариошкин С.Н.* Современные представления об этиологии болезни Паркинсона. Неврол. журн. 2015; 20(4): 4–13.

*Иллариошкин С.Н.* Терапевтическое воздействие электричества на головной мозг: от первых шагов времен античности до современных методов глубокой электростимуляции мозга. Бюллетень Национального общества по изучению болезни Паркинсона и расстройств движений. 2013а; 1: 26–32.

*Иллариошкин С.Н.* Терапия паркинсонизма: возможности и перспективы. Consilium Medicum. Неврол. и ревматол. 2009; 1: 35–40.

*Иллариошкин С.Н.* Этапы прогрессирования болезни Паркинсона. Что определяет выживаемость и прогноз? Вестник Российской Военно-мед. акад. 2013б; 4(прил. 2): 11–13.

*Иллариошкин С.Н., Иванова-Смоленская И.А.* Дрожательные гиперкинезы. Руководство для врачей. М.: Атмосфера; 2011. 360 с.

*Иллариошкин С.Н., Левин О.С.* Руководство по диагностике и лечению болезни Паркинсона. М.: МЕДпресс-информ; 2017. 336 с.

*Иллариошкин С.Н., Левин О.С., Федотова Е.Ю., Колоколов О.В.* Паркинсонизм и черная субстанция. М.: Издательское предприятие «Атмосфера»; 2019. 304 с.

*Кандель Э.И.* Функциональная и стереотаксическая нейрохирургия. М.: Медицина; 1981. 368 с.

*Кандель Э.И., Войтына С.В.* Деформирующая мышечная (торсионная) дистония. Клиника и хирургическое лечение. М.: Медицина; 1971. 184 с.

*Левин О.С., Иллариошкин С.Н., Голубев В.Л.* Экстрапирамидные синдромы. Руководство для врачей. М.: МЕДпресс-информ; 2022. 772 с.

*Лихтерман Л.Б.* Воспоминания об Эдуарде Израилевиче Канделе (1923–1990). Журн. «Вопросы нейрохирургии» им. Н.Н. Бурденко. 2013; 77(4): 69–72.

*Михайлов В.А., Дружинин А.К., Шова Н.И., Корсакова Е.А.* Эпилепсия с геластическими приступами и гамартома гипоталамуса (описание редкого клинического наблюдения). Эпилепсия и пароксизм. состояния. 2017; 9(3): 64–72.

*Низаметдинова Д.М., Тюрников В.М., Федоренко И.И. и др.* Микроэлектродная регистрация нейрональной активности в хирургии болезни Паркинсона. Анн. клин. и эксперимент. неврол. 2016; 10(2): 42–45.

*Резников И.И., Федорова В.Н., Фаустов Е.В.* Физические основы использования ультразвука в медицине. М.: РНИМУ им. Н.И. Пирогова; 2015. 97 с.

*Розенберг Л.Д.* Звуковые фокусирующие системы. М.: АН СССР; 1949. 112 с.

*Розенберг Л.Д.* Фокусирующие излучатели ультразвука. В кн.: Физика и техника мощного ультразвука. М.: Наука; 1967: 149–206.

Росздравнадзор (Федеральная служба по контролю в сфере здравоохранения). Регистрационное удостоверение от 13 февраля 2017 года на медицинское изделие № РЗН 2017/5378. Система для неинвазивной абляции фокусированным ультразвуком ExAblate с принадлежностями.

*Саветеев А.Н., Голанов А.В., Саушев Д.А. и др.* Стереотаксическая радиохирургия при эпилепсии у пациентов с гамартомой гипоталамуса. Журн. «Вопросы нейрохирургии» им. Н.Н. Бурденко. 2022; 86(4): 14–24.

*Сунгуров Е.Б.* Отдаленные результаты хирургического лечения стереотаксическим криогенным методом больных торсионной дистонией и сходных синдромов: Автореф. дис. ... канд. мед. наук. М., 1997. 27 с.

*Тюрников В.М.* Нейрохирургическое лечение синдрома Туретта: когда необходимо оперативное вмешательство. Практический психиатр и психолог. 2022; 11: 1–6.

*Тюрников В.М., Гуца А.О.* Высокоинтенсивный фокусированный ультразвук в функциональной нейрохирургии. Анн. клин. и эксперимент. неврол. 2016; 10(4): 52–57.

*Тюрников В.М., Маркова Е.Д., Добжанский Н.В.* Хирургическое лечение краниальной и цервикальной дистонии. Атмосфера. Нервные болезни. 2007; 3: 28–32.

Холявин А.И. Технические, клинические и экономические аспекты лечебного транскраниального фокусированного ультразвука. Бюллетень Национального общества по изучению болезни Паркинсона и расстройств движений. 2019; 1: 17–24.

Шабалов В.А. Нейрохирургическое лечение экстрапирамидных расстройств. В кн.: Экстрапирамидные расстройства. Руководство по диагностике и лечению. Под ред. Штока В.Н., Ивановой-Смоленской И.А., Левина О.С. М.: МЕДпресс-информ; 2002: 552–566.

Шабалов В.А., Исагулян Э.Д. Хроническая электростимуляция в лечении невропатических болевых синдромов. Критерии длительной эффективности. Нервные болезни. 2010; 4: 2–9.

Шабалов В.А., Томский А.А. Хирургическое лечение болезни Паркинсона (часть 1). Нейрохирургия. 2003; 3: 5–16.

Abe K., Horisawa S., Yamaguchi T. et al. Focused ultrasound thalamotomy for refractory essential tremor: a Japanese multicenter single-arm study. Neurosurgery. 2021 Mar; 88(4): 751–757.

Abrahamo A., Meng Y., Llinas M. et al. First-in-human trial of blood-brain barrier opening in amyotrophic lateral sclerosis using MR-guided focused ultrasound. Nat. Commun. 2019 Sep; 10(1): 4373.

Akbostanci M.C., Slavin K.V., Burchiel K.J. Stereotactic ventral intermedial thalamotomy for the treatment of essential tremor: results of a series of 37 patients. Stereotact. Funct. Neurosurg. 1999; 72(2-4): 174–177.

Albanese A., Bhatia K., Bressman S.B. et al. Phenomenology and classification of dystonia: a consensus update. Mov. Disord. 2013 Jun; 28(7): 863–873.

Albanese A., Di Giovanni M., Lalli S. Dystonia: diagnosis and management. Eur. J. Neurol. 2019 Jan; 26(1): 5–17.

Al-Bataineh O., Jenne J., Huber P. Clinical and future applications of high intensity focused ultrasound in cancer. Cancer Treat. Rev. 2012 Aug; 38(5): 346–353.

Albin R.L., Young A.B., Penney J.B. The functional anatomy of basal ganglia disorders. Trends Neurosci. 1989 Oct; 12(10): 366–375.

Allam A.K., Larkin M.B., McGinnis J.P., Viswanathan A. Neuroablative central lateral thalamotomy for chronic neuropathic pain. Front. Pain Res. (Lausanne). 2022 Sep; 3: 999891.

Alshaikh J., Fishman P.S. Revisiting bilateral thalamotomy for tremor. Clin. Neurol. Neurosurg. 2017 Jul; 158: 103–107.

Alterman R., Snyder B. Deep brain stimulation for torsion dystonia. Acta Neurochir. Suppl. 2007; 97(Pt. 2): 191–199.

Altinel Y., Alkhalfan F., Qiao N., Velimirovic M. Outcomes in lesion surgery versus deep brain stimulation in patients with tremor: a systematic review and meta-analysis. World Neurosurg. 2019 Mar; 123: 443–452.e8.

Alvarez-Garjoto J.A., Albiach V.J., Vila M.M. et al. Precocious puberty and hypothalamic hamartoma with total recovery after surgical treatment: case report. J. Neurosurg. 1983 Apr; 58(4): 583–585.

Artusi C.A., Dwivedi A., Romagnolo A. et al. Differential response to pallidal deep brain stimulation among monogenic dystonias: systematic review and meta-analysis. J. Neurol. Neurosurg. Psychiatry. 2020 Apr; 91(4): 426–433.

Aufenberg C., Sarnthein J., Morel A. et al. A revival of Spiegel's camptomy: long term results of the stereotactic pallidothalamic tractotomy against the parkinsonian thalamocortical dysrhythmia. Thalamus Relat. Syst. 2005 Jun; 3(2): 121–132.

Baek H., Lockwood D., Mason E.J. et al. Clinical intervention using focused ultrasound (FUS) stimulation of the brain in diverse neurological disorders. Front. Neurol. 2022 May; 13: 880814.

Bain P.G., Findley L.J., Thompson P.D. et al. A study of hereditary essential tremor. Brain. 1994 Aug; 117(Pt. 4): 805–824.

Bechtereva N.P., Bondartchuk A.N., Smirnov V.M. et al. Method of electrostimulation of the deep brain structures in treatment of some chronic diseases. Confin. Neurol. 1975; 37(1-3): 136–140.

- Beck A.T., Ward C.H., Mendelson M. et al. An inventory for measuring depression. *Arch. Gen. Psychiatry.* 1961 Jun; 4: 561–571.
- Beggs J., Nakada S., Fenoglio K. et al. Hypothalamic hamartomas associated with epilepsy: ultrastructural features. *J. Neuropathol. Exp. Neurol.* 2008 Jul; 67(7): 657–668.
- Benabid A.L., Koudsié A., Benazzouz A. et al. Subthalamic stimulation for Parkinson's disease. *Arch. Med. Res.* 2000 May-Jun; 31(3): 282–289.
- Benabid A.L., Pollak P., Gervason C. et al. Long-term suppression of tremor by chronic stimulation of the ventral intermediate thalamic nucleus. *Lancet.* 1991 Feb; 337(8738): 403–406.
- Benabid A.L., Pollak P., Louveau A. et al. Combined (thalamotomy and stimulation) stereotactic surgery of the VIM thalamic nucleus for bilateral Parkinson disease. *Appl. Neurophysiol.* 1987; 50(1-6): 344–346.
- Benabid A.L., Pollak P., Seigneuret E. et al. Chronic VIM thalamic stimulation in Parkinson's disease, essential tremor and extra-pyramidal dyskinesias. *Acta Neurochir. Suppl.* 1993; 58: 39–44.
- Berardelli A., Wenning G.K., Antonini A. et al. EFNS/MDS-ES recommendations for the diagnosis of Parkinson's disease. *Eur. J. Neurol.* 2013 Jan; 20(1): 16–34.
- Berendse H.W., van Laar T. Tremor. In: Parkinsonism and related disorders. Wolters E.Ch., van Laar T., Berendse H.W., editors. Amsterdam: VU University Press; 2007: 309–325.
- Bernasconi A., Cendes F., Theodore W.H. et al. Recommendations for the use of structural magnetic resonance imaging in the care of patients with epilepsy: a consensus report from the International League Against Epilepsy Neuroimaging Task Force. *Epilepsia.* 2019; 60(6): 1054–1068.
- Bhatia K.P., Bain P., Bajaj N. et al. Consensus statement on the classification of tremors from the task force on tremor of the International Parkinson and Movement Disorder Society. *Mov. Disord.* 2018 Jan; 33(1): 75–87.
- Blahak C., Wolf M.E., Saryyeva A. et al. Improvement of head and neck range of motion induced by chronic pallidal deep brain stimulation for cervical dystonia. *J. Neural Transm. (Vienna).* 2021 Aug; 128(8): 1205–1213.
- Blomstedt P., Hariz G.M., Hariz M.I. Pallidotomy versus pallidal stimulation. *Parkinsonism Relat. Disord.* 2006 Jun; 12(5): 296–301.
- Bond A.E., Elias W.J. Predicting lesion size during focused ultrasound thalamotomy: a review of 63 lesions over 3 clinical trials. *Neurosurg. Focus.* 2018 Feb; 44(2): E5.
- Bond A.E., Shah B.B., Huss D.S. et al. Safety and efficacy of focused ultrasound thalamotomy for patients with medication-refractory, tremor-dominant Parkinson disease: a randomized clinical trial. *JAMA Neurol.* 2017 Dec; 74(12): 1412–1418.
- Bouhassira D., Lantéri-Minet M., Attal N. et al. Prevalence of chronic pain with neuropathic characteristics in the general population. *Pain.* 2008 Jun; 136(3): 380–387.
- Braak H., Del Tredici K., Rüb U. et al. Staging of brain pathology related to sporadic Parkinson's disease. *Neurobiol. Aging.* 2003 Mar-Apr; 24(2): 197–211.
- Bradley W.G. Jr. MR-guided focused ultrasound: a potentially disruptive technology. *J. Am. Coll. Radiol.* 2009 Jul; 6(7): 510–513.
- Brahmandam A., Chan S.M., Dardik A. et al. A narrative review on the application of high-intensity focused ultrasound for the treatment of occlusive and thrombotic arterial disease. *JVS Vasc. Sci.* 2022 Sep; 3: 292–305.
- Brice J., McLellan L. Suppression of intention tremor by contingent deep-brain stimulation. *Lancet.* 1980 Jun; 315(8180): 1221–1222.
- Broering D.C., Berardi G., El Sheikh Y. et al. Learning curve under proctorship of pure laparoscopic living donor left lateral sectionectomy for pediatric transplantation. *Ann. Surg.* 2020 Mar; 271(3): 542–548.

- Brown M.R., Farquhar-Smith P., Williams J.E. et al.* The use of high-intensity focused ultrasound as a novel treatment for painful conditions – a description and narrative review of the literature. *Br. J. Anaesth.* 2015 Oct; 115(4): 520–530.
- Bruno F., Catalucci A., Varrassi M. et al.* Comparative evaluation of tractography-based direct targeting and atlas-based indirect targeting of the ventral intermediate (Vim) nucleus in MRgFUS thalamotomy. *Sci. Rep.* 2021 Jan; 11(1): 13538.
- Burke R.E., Fahn S., Marsden C.D. et al.* Validity and reliability of a rating scale for the primary torsion dystonias. *Neurology.* 1985 Jan; 35(1): 73–77.
- Butler A.G., Duffey P.O., Hawthorne M.R. et al.* The socioeconomic implications of dystonia. *Adv. Neurol.* 1998; 78: 349–358.
- Buzaev I.V., Galimova R.M., Nabiullina D.I. et al.* Magnetic resonance imaging-guided focused ultrasound thalamotomy launch with remote telemedicine international proctorship. *Chronic Dis. Transl. Med.* 2023 Sep; 10(1): 40–50.
- Cacho-Asenjo E., Honorato-Cia C., Nuñez-Cordoba J.M. et al.* Factors associated with headache and nausea during magnetic resonance guided focused ultrasound for tremor. *Mov. Disord. Clin. Pract.* 2021 Apr; 8(5): 701–708.
- Castro L.H., Ferreira L.K., Teles L.R. et al.* Epilepsy syndromes associated with hypothalamic hamartomas. *Seizure.* 2007 Jan; 16(1): 50–58.
- Chahine L.M., Merchant K., Siderowf A. et al.* Proposal for a biologic staging system of Parkinson's disease. *J. Parkinsons Dis.* 2023; 13(3): 297–309.
- Chang J.W., Park C.K., Lipsman N. et al.* A prospective trial of magnetic resonance-guided focused ultrasound thalamotomy for essential tremor: results at the 2-year follow-up. *Ann. Neurol.* 2018 Jan; 83(1): 107–114.
- Chang W.S., Jung H.H., Kweon E.J. et al.* Unilateral magnetic resonance guided focused ultrasound thalamotomy for essential tremor: practices and clinicoradiological outcomes. *J. Neurol. Neurosurg. Psychiatry.* 2015 Mar; 86(3): 257–264.
- Chatterjee A., Jurewicz E.C., Applegate L.M., Louis E.D.* Personality in essential tremor: further evidence of non-motor manifestations of the disease. *J. Neurol. Neurosurg. Psychiatry.* 2004 Jul; 75(7): 958–961.
- Chau M.T., Todd G., Wilcox R. et al.* Diagnostic accuracy of the appearance of nigrosome-1 on magnetic resonance imaging in Parkinson's disease: a systematic review and meta-analysis. *Parkinsonism Relat. Disord.* 2020 Sep; 78: 12–20.
- Chaudhuri K.R., Healy D.G., Schapira A.H.* Non-motor symptoms of Parkinson's disease: diagnosis and management. *Lancet. Neurol.* 2006 Mar; 5(3): 235–245.
- Chen C.J., Hsu H.C., Chung W.S., Yu H.J.* Clinical experience with ultrasound-based real-time tracking lithotripsy in the single renal stone treatment. *J. Endourol.* 2009 Nov; 23(11): 1811–1815.
- Clement G.T., White J., Hynynen K.* Investigation of a large-area phased array for focused ultrasound surgery through the skull. *Phys. Med. Biol.* 2000 Apr; 45(4): 1071–1083.
- Cohen N.T., Cross J.H., Arzimanoglou A. et al.; Hypothalamic Hamartoma Writing Group.* Hypothalamic hamartomas: evolving understanding and management. *Neurology.* 2021 Nov; 97(18): 864–873.
- Cohen Z.R., Zauber mann J., Harnof S. et al.* Magnetic resonance imaging-guided focused ultrasound for thermal ablation in the brain: a feasibility study in a swine model. *Neurosurgery.* 2007 Apr; 60(4): 593–600.
- Coleman C., Martin I.* Unraveling Parkinson's disease neurodegeneration: does aging hold the clues? *J. Parkinsons Dis.* 2022; 12(8): 2321–2338.
- Collis J., Manasseh R., Petar L. et al.* Cavitation microstreaming and stress fields created by microbubbles. *Ultrasonics.* 2010 Feb; 50(2): 273–279.
- Comella C.L., Fox S.H., Bhatia K.P. et al.* Reliability of the severity Subscale of the revised Toronto Spasmodic Torticollis Rating Scale (TWSTRS-2) (S15.001). *Neurology* 2015; 84(14 Suppl.): S15.001.

---

*Comella C.L., Leurgans S., Wu J. et al.* Rating scales for dystonia: a multicenter assessment. *Mov. Disord.* 2003 Mar; 18(3): 303–312.

*Consky E.S., Lang A.E.* Clinical assessments of patients with cervical dystonia. In: Jankovic J., Hallett M., editors. *Therapy with Botulinum Toxin*. New York: Marcel Dekker; 1994: 211–237.

*Cooper I.S.* Involuntary movement disorders. New York: Hoeber Medical Division, Harper & Row; 1969. 410 p.

*Cooper I.S.* Ligation of the anterior choroidal artery for involuntary movements; parkinsonism. *Psychiatr. Q.* 1953 Apr; 27(2): 317–319.

*Cooper I.S., Upton A.R.* Therapeutic implications of modulation of metabolism and functional activity of cerebral cortex by chronic stimulation of cerebellum and thalamus. *Biol. Psychiatry.* 1985 Jul; 20(7): 811–813.

*Cooper I.S., Upton A.R., Amin I.* Chronic cerebellar stimulation (CCS) and deep brain stimulation (DBS) in involuntary movement disorders. *Appl. Neurophysiol.* 1982; 45(3): 209–217.

*Coubes P., Echenne B., Roubertie A. et al.* [Treatment of early-onset generalized dystonia by chronic bilateral stimulation of the internal globus pallidus. Apropos of a case.] (In French). *Neurochirurgie.* 1999 May; 45(2): 139–144.

*Dalecki D.* Mechanical bioeffects of ultrasound. *Annu. Rev. Biomed. Eng.* 2004; 6: 229–248.

*Darrow D.P.* Focused ultrasound for neuromodulation. *Neurotherapeutics.* 2019 Jan; 16(1): 88–99.

*Davidson B., Hamani C., Huang Y. et al.* Magnetic resonance-guided focused ultrasound capsulotomy for treatment-resistant psychiatric disorders. *Oper. Neurosurg.* 2020a Nov; 19(6): 741–749.

*Davidson B., Hamani C., Rabin J.S. et al.* Magnetic resonance-guided focused ultrasound capsulotomy for refractory obsessive compulsive disorder and major depressive disorder: clinical and imaging results from two phase I trials. *Mol. Psychiatry.* 2020b Sep; 25(9): 1946–1957.

*Dayton P., Klibanov A., Brandenburger G., Ferrara K.* Acoustic radiation force *in vivo*: a mechanism to assist targeting of microbubbles. *Ultrasound Med. Biol.* 1999 Oct; 25(8): 1195–1201.

*De Bie R.M., Schuurman P.R., Esselink R.A. et al.* Bilateral pallidotomy in Parkinson's disease: a retrospective study. *Mov. Disord.* 2002 May; 17(3): 533–538.

*de Natale E.R., Niccolini F., Wilson H., Politis M.* Molecular imaging of the dopaminergic system in idiopathic Parkinson's disease. *Int. Rev. Neurobiol.* 2018; 141: 131–172.

*Delalande O., Fohlen M.* Disconnecting surgical treatment of hypothalamic hamartoma in children and adults with refractory epilepsy and proposal of a new classification. *Neurol. Med. Chir.* 2003 Feb; 43(2): 61–68.

*Deng C.X.* Targeted drug delivery across the blood brain barrier using ultrasound technique. *Ther. Deliv.* 2010 Dec; 1(6): 819–848.

*Dogu O., Sevim S., Camdeviren H. et al.* Prevalence of essential tremor: door-to-door neurologic exams in Mersin Province, Turkey. *Neurology.* 2003 Dec; 61(12): 1804–1806.

*Domingo A., Yadav R., Ozelius L.J.* Isolated dystonia: clinical and genetic updates. *J. Neural Transm. (Vienna).* 2021 Apr; 128(4): 405–416.

*Driver J.A., Logroscino G., Gaziano J.M., Kurth T.* Incidence and remaining lifetime risk of Parkinson disease in advanced age. *Neurology.* 2009 Feb; 72(5): 432–438.

*Dubey S., Heinen S., Krantic S. et al.* Clinically approved IVIg delivered to the hippocampus with focused ultrasound promotes neurogenesis in a model of Alzheimer's disease. *Proc. Natl. Acad. Sci. USA.* 2020 Dec; 117(51): 32691–32700.

*Duma C., Jacques D., Kopyov O. et al.* Gamma knife radiosurgery for thalamotomy in parkinsonian tremor: a five-year experience. *J. Neurosurg.* 1998 Jun; 88(6): 1044–1049.

*Durif F., Lemaire J.J., Debilly B., Dordain G.* Long-term follow-up of globus pallidus chronic stimulation in advanced Parkinson's disease. *Mov. Disord.* 2002 Jul; 17(4): 803–807.

- Eisen A., Braak H., Del Tredici K. et al. Cortical influences drive amyotrophic lateral sclerosis. *J. Neurol. Neurosurg. Psychiatry*. 2017 Nov; 88(11): 917–924.
- Eisenberg H.M., Krishna V., Elias W.J. et al. MR-guided focused ultrasound pallidotomy for Parkinson's disease: safety and feasibility. *J. Neurosurg*. 2020 Nov; 135(3): 792–798.
- Elble R.J. Do we belittle essential tremor by calling it a syndrome rather than a disease? No. *Front. Neurol*. 2020 Sep; 11: 586606.
- Elble R.J. The Essential Tremor Rating Assessment Scale. *J. Neurol. Neuromed*. 2016; 1(4): 34–38.
- Elias W.J., Huss D., Voss T. et al. A pilot study of focused ultrasound thalamotomy for essential tremor. *N. Engl. J. Med*. 2013 Aug; 369(7): 640–648.
- Elias W.J., Lipsman N., Ondo W.G. et al. A randomized trial of focused ultrasound thalamotomy for essential tremor. *N. Engl. J. Med*. 2016 Aug; 375(8): 730–739.
- Engel J. Jr. Outcome with respect to epileptic seizures. In: *Surgical treatment of the epilepsies*. Engel J. Jr, editor. 2nd ed. New York: Raven Press; 1993: 609–621.
- Espay A.J., Hausdorff J.M., Sanchez-Ferro Á. et al. A roadmap for implementation of patient-centered digital outcome measures in Parkinson's disease obtained using mobile health technologies. *Mov. Disord*. 2019 May; 34(5): 657–663.
- Fahn S., Tolosa E., Conception M. Clinical rating scale for tremor. In: *Parkinson's disease and movement disorders*. 2nd ed. Jankovich J., Tolosa E., editors. Baltimore, MD: Williams & Wilkins; 1993: 271–280.
- Fahn S., Tolosa E., Marin C. Clinical rating scale for tremor. In: *Parkinson's disease and movement disorders*. Jankovich J., Tolosa E., editors. Baltimore, MD: Urban & Schwarzenberg; 1988: 225–234.
- Fasano A., Deuschl G. Therapeutic advances in tremor. *Mov. Disord*. 2015; 30(11): 1557–1565.
- Fasano A., De Vloot P., Llinas M. et al. Magnetic resonance imaging-guided focused ultrasound thalamotomy in Parkinson tremor: reoperation after benefit decay. *Mov. Disord*. 2018 May; 33(5): 848–849.
- FDA News Release. U.S. Food and Drug Administration. FDA approves first MRI-guided focused ultrasound device to treat essential tremor. Posted; 2016 Jul 11. Available from: <https://www.fda.gov/news-events/press-announcements/fda-approves-first-mri-guided-focused-ultrasound-treat-essential-tremor> Accessed 2024 Mar 21.
- Fenoy A.J., Goetz L., Chabardès S., Xia Y. Deep brain stimulation: are astrocytes a key driver behind the scene? *CNS Neurosci. Ther*. 2014 Mar; 20(3): 191–201.
- Ferguson T., Frydman G. The first generation of e-patients. *BMJ*. 2004 May; 328(7449): 1148–1149.
- Ferreira J.J., Mestre T.A., Lyons K.E. et al. MDS evidence-based review of treatments for essential tremor. *Mov. Disord*. 2019 Jul; 34(7): 950–958.
- Fine J., Duff J., Chen R. et al. Long-term follow-up of unilateral pallidotomy in advanced Parkinson's disease. *N. Engl. J. Med*. 2000 Jun; 342(23): 1708–1714.
- Fishman P.S., Frenkel V. Treatment of movement disorders with focused ultrasound. *J. Cent. Nerv. Syst. Dis*. 2017 Jun; 9: 1179573517705670.
- Foley J.L., Eames M., Snell J. et al. Image-guided focused ultrasound: state of the technology and the challenges that lie ahead. *Imag. Med*. 2013; 5(4): 357.
- Fox S.H., Katzenschlager R., Lim S.Y. et al.; *Disorder Society Evidence-Based Medicine Committee*. International Parkinson and Movement Disorder Society evidence-based medicine review: update on treatments for the motor symptoms of Parkinson's disease. *Mov. Disord*. 2018 Aug; 33(8): 1248–1266.
- Freeman W.J., Watts J.W. Retrograde degeneration of the thalamus following prefrontal lobotomy. *J. Compar. Neurol*. 1947 Feb; 86(1): 65–93.

- Freund B., Sutter R., Kaplan P.W. Lance–Adams syndrome in the pretargeted temperature management era: a case report and systematic review. *Clin. EEG Neurosci.* 2017 Mar; 48(2): 130–138.
- Frucht S.J., Leurgans S.E., Hallett M., Fahn S. The Unified Myoclonus Rating Scale. *Adv. Neurol.* 2002; 89: 361–376.
- Fry F.J. Precision high intensity focusing ultrasonic machines for surgery. *Am. J. Phys. Med. Rehabil.* 1958 Jun; 37(3): 152–156.
- Fry W.J., Meyers R. Ultrasonic method of modifying brain structures. *Confin. Neurol.* 1962; 22: 315–327.
- Fukutome K., Hirabayashi H., Osakada Y. et al. Bilateral magnetic resonance imaging-guided focused ultrasound thalamotomy for essential tremor. *Stereotact. Funct. Neurosurg.* 2022; 100(1): 44–52.
- Gagliardo C., Midiri M., Cannella R. et al. Transcranial magnetic resonance imaging-guided focused ultrasound treatment at 1.5-T: a retrospective study on treatment- and patient-related parameters obtained from 52 procedures. *Neuroradiol. J.* 2019; 32(2): 132–138.
- Galimova R., Kachemaeva O., Akhmadeeva G. et al. First in history MRgFUS treatment procedures remote start and successful treatment of PD patients due to COVID-19 epidemic. *Mov. Disord.* 2021; 36(Suppl. 1): S542.
- Galimova R., Sidorova Yu., Akhmadeeva G. et al. Transcranial magnetic resonance-guided focused ultrasound in Lance–Adams syndrome. *Mov. Disord.* 2022; 37(Suppl. 1): 281.
- Gallay M.N., Moser D., Jeanmonod D. MR-guided focused ultrasound central lateral thalamotomy for trigeminal neuralgia. Single center experience. *Front. Neurol.* 2020a Apr; 11: 271.
- Gallay M.N., Moser D., Jeanmonod D. Safety and accuracy of incisionless transcranial MR-guided focused ultrasound functional neurosurgery: single-center experience with 253 targets in 180 treatments. *J. Neurosurg.* 2018 May; 130(4): 1234–1243.
- Gallay M.N., Moser D., Magara A.E. et al. Bilateral MR-guided focused ultrasound pallidothalamic tractotomy for Parkinson’s disease with 1-year follow-up. *Front. Neurol.* 2021 Feb; 12: 601153.
- Gallay M.N., Moser D., Rossi F. et al. Incisionless transcranial MR-guided focused ultrasound in essential tremor: cerebellothalamic tractotomy. *J. Ther. Ultrasound.* 2016 Feb; 4: 5.
- Gallay M.N., Moser D., Rossi F. et al. MRgFUS pallidothalamic tractotomy for chronic therapy-resistant Parkinson’s disease in 51 consecutive patients: single center experience. *Front. Surg.* 2020b; 6: 76.
- Gervais-Bernard H., Xie-Brustolin J., Mertens P. et al. Bilateral subthalamic nucleus stimulation in advanced Parkinson’s disease: five-year follow-up. *J. Neurol.* 2009 Feb; 256(2): 225–233.
- Geyer L.H., Bressman S.B. The diagnosis of dystonia. *Lancet. Neurol.* 2006 Sep; 5(9): 780–790.
- Giordano M., Caccavella V.M., Zaed I. et al. Comparison between deep brain stimulation and magnetic resonance-guided focused ultrasound in the treatment of essential tremor: a systematic review and pooled analysis of functional outcomes. *J. Neurol. Neurosurg. Psychiatry.* 2020 Dec; 91(12): 1270–1278.
- Godinho F., Magnin M., Filho P.T. et al. Stereotactic lesion in the Forel’s field H: a 2-years prospective open-label study on motor and nonmotor symptoms, neuropsychological functions, and quality of life in Parkinson disease. *Neurosurgery.* 2019 Oct; 85(4): E650–E659.
- Goetz C.G., Poewe W., Rascol O. et al. Movement Disorder Society Task Force on Rating Scales for Parkinson’s Disease. Movement Disorder Society Task Force report on the Hoehn and Yahr Staging Scale: status and recommendation. *Mov. Disord.* 2004 Sep; 19(9): 1020–1028.
- Goetz C.G., Tilley B.C., Shaftman S.R. et al. Movement Disorder Society-sponsored revision of the Unified Parkinson’s Disease Rating Scale (MDS-UPDRS): scale presentation and clinimetric testing results. *Mov. Disord.* 2008 Nov; 23(15): 2129–2170.
- Goldman M.S., Kelly P.J. The surgical treatment of tremor disorders. In: *Handbook of tremor disorders.* Findley L.J., Koller W.C., editors. New York: Marcel Dekker; 1995: 521–562.



- Golfrè Andreasi N, Braccia A, Levi V.* The optimal targeting for focused ultrasound thalamotomy differs between dystonic and essential tremor: a 12-month prospective pilot study. *Mov. Disord. Clin. Pract.* 2024 Jan; 11(1): 69–75.
- Goodman S.H., Wilkison S., Overman J. et al.* Lesion volume and clinical outcome in stereotactic pallidotomy and thalamotomy. *Stereotact. Funct. Neurosurg.* 1998; 71(4): 164–172.
- Gruetzmacher J.* Piezoelektrischer kristall mit ultraschallkonvergenz. *Zeitschr. Physik.* 1935; 96(5-6): 342–349.
- Hacker M.L., DeLong M.R., Turchan M. et al.* Effects of deep brain stimulation on rest tremor progression in early-stage Parkinson disease. *Neurology.* 2018 Jul; 91(5): e463–e471.
- Hacker M.L., Turchan M., Heusinkveld L.E. et al.* Deep brain stimulation in early-stage Parkinson disease: five-year outcomes. *Neurology.* 2020 Jul; 95(4): e393–e401.
- Halpern C.H., Santini V., Lipsman N. et al.* Three-year follow-up of prospective trial of focused ultrasound thalamotomy for essential tremor. *Neurology.* 2019 Dec; 93(24): e2284–e2293.
- Hardy C.J., Cline H.E., Watkins R.D.* One-dimensional NMR thermal mapping of focused ultrasound surgery. *J. Comput. Assist. Tomogr.* 1994 May-Jun; 18(3): 476–483.
- Hariz G.M., Lindberg M., Bergenheim A.T.* Impact of thalamic deep brain stimulation on disability and health-related quality of life in patients with essential tremor. *J. Neurol. Neurosurg. Psychiatry.* 2002 Jan; 72(1): 47–52.
- Hariz M.I., Bergenheim A.T.* A 10-year follow-up review of patients who underwent Leksell's posteroventral pallidotomy for Parkinson disease. *J. Neurosurg.* 2001 Apr; 94(4): 552–558.
- Hariz M.I., Robertson M.M.* Gilles de la Tourette syndrome and deep brain stimulation. *Eur. J. Neurosci.* 2010 Oct; 32(7): 1128–1134.
- Hassler R.* Extrapyramidal-motorische Syndrome und Erkrankungen. In: *Handbuch der inneren Medizin. Neurologie III.* Berlin–Göttingen–Heidelberg: Springer; 1953. 676 p.
- Health Quality Ontario.* Magnetic resonance-guided focused ultrasound neurosurgery for essential tremor: a health technology assessment. *Ont. Health Technol. Assess. Ser.* 2018 May; 18(4): 1–141.
- Heras-Garvin A., Stefanova N.* From synaptic protein to prion: the long and controversial journey of  $\alpha$ -synuclein. *Front. Synaptic Neurosci.* 2020 Sep; 12: 584536.
- Hoehn M.M., Yahr M.D.* Parkinsonism: onset, progression, and mortality. *Neurology.* 1967 May; 17(5): 427–442.
- Horisawa S., Fukui A., Yamahata H. et al.* Unilateral pallidothalamic tractotomy for akinetic-rigid Parkinson's disease: a prospective open-label study. *J. Neurosurg.* 2021a Jan; 135(3): 799–805.
- Horisawa S., Yamaguchi T., Abe K. et al.* A single case of MRI-guided focused ultrasound ventro-oral thalamotomy for musician's dystonia. *J. Neurosurg.* 2018 Sep; 131(2): 384–386.
- Horisawa S., Yamaguchi T., Abe K. et al.* Magnetic resonance-guided focused ultrasound thalamotomy for focal hand dystonia: a pilot study. *Mov. Disord.* 2021b Aug; 36(8): 1955–1959.
- Horsley V., Clarke R.H.* The structure and functions of the cerebellum examined by a new method. *Brain.* 1908 May; 31(1): 45–124.
- Huang H., Yang X., Zhao Q. et al.* Prevalence and risk factors of depression and anxiety in essential tremor patients: a cross-sectional study in Southwest China. *Front. Neurol.* 2019 Nov; 10: 1194.
- Huss D.S., Dallapiazza R.F., Shah B.B. et al.* Functional assessment and quality of life in essential tremor with bilateral or unilateral DBS and focused ultrasound thalamotomy. *Mov. Disord.* 2015 Dec; 30(14): 1937–1943.
- Hynynen K., Damianou C., Darkazanli A. et al.* The feasibility of using MRI to monitor and guide noninvasive ultrasound surgery. *Ultrasound Med. Biol.* 1993; 19(1): 91–92.
- Hynynen K., Freund W.R., Cline H.E. et al.* A clinical, noninvasive, MR imaging-monitored ultrasound surgery method. *Radiographics.* 1996 Jan; 16(1): 185–195.

- Hynynen K., Jolesz F.A.* Demonstration of potential noninvasive ultrasound brain therapy through an intact skull. *Ultrasound Med. Biol.* 1998 Feb; 24(2): 275–283.
- Hynynen K., McDannold N., Clement G. et al.* Pre-clinical testing of a phased array ultrasound system for MRI-guided noninvasive surgery of the brain – a primate study. *Eur. J. Radiol.* 2006 Aug; 59(2): 149–156.
- Iacopino D.G., Gagiardo C., Giugno A. et al.* Preliminary experience with a transcranial magnetic resonance-guided focused ultrasound surgery system integrated with a 1.5-T MRI unit in a series of patients with essential tremor and Parkinson's disease. *Neurosurg. Focus.* 2018 Feb; 44(2): E7.
- Intemann P.M., Masterman D., Subramanian I. et al.* Staged bilateral pallidotomy for treatment of Parkinson disease. *J. Neurosurg.* 2001 Mar; 94(3): 437–444.
- Iorio-Morin C., Yamamoto K., Sarica C. et al.* Bilateral focused ultrasound thalamotomy for essential tremor (BEST-FUS phase 2 trial). *Mov. Disord.* 2021 Nov; 36(11): 2653–2662.
- Jackson L.M., Kaufmann T.J., Lehman V.T. et al.* Clinical characteristics of patients with gait instability after MR-guided focused ultrasound thalamotomy. *Tremor Other Hyperkinet. Mov. (NY).* 2021; 11: 41.
- Jagannathan J., Sanghvi N.T., Crum L.A. et al.* High-intensity focused ultrasound surgery of the brain: part 1 – a historical perspective with modern applications. *Neurosurgery.* 2009 Feb; 64(2): 201–210; discussion 210–211.
- Jameel A., Gedroyc W., Nandi D. et al.* Double lesion MRgFUS treatment of essential tremor targeting the thalamus and posterior sub-thalamic area: preliminary study with two-year follow-up. *Br. J. Neurosurg.* 2022; 36(2): 241–250.
- Jamora R.D.G., Chang W.C., Taira T.* Transcranial magnetic resonance-guided focused ultrasound in X-linked dystonia-parkinsonism. *Life.* 2021 Apr; 11(5): 392.
- Jang H.J., Lee J.Y., Lee D.H. et al.* Current and future clinical applications of High-Intensity Focused Ultrasound (HIFU) for pancreatic cancer. *Gut Liver.* 2010 Sep; 4(Suppl. 1): S57–S61.
- Jeanmonod D., Magnin M., Morel A.* Chronic neurogenic pain and the medial thalamotomy. *Schweiz. Rundsc. Med. Praxis.* 1994 Jun; 83(23): 702–707.
- Jeanmonod D., Werner B., Morel A. et al.* Transcranial magnetic resonance imaging-guided focused ultrasound: noninvasive central lateral thalamotomy for chronic neuropathic pain. *Neurosurg. Focus.* 2012 Jan; 32(1): E1.
- Jenner P., Morris H.R., Robbins T.W. et al.* Parkinson's disease – the debate on the clinical phenomenology, aetiology, pathology and pathogenesis. *J. Parkinsons Dis.* 2013; 3(1): 1–11.
- Johns M.W.* A new method for measuring daytime sleepiness: the Epworth Sleepiness Scale. *Sleep.* 1991 Dec; 14(6): 540–545.
- Jung H.H., Kim S.J., Roh D. et al.* Bilateral thermal capsulotomy with MR-guided focused ultrasound for patients with treatment-refractory obsessive-compulsive disorder: a proof-of-concept study. *Mol. Psychiatry.* 2015 Oct; 20(10): 1205–1211.
- Jung N.Y., Chang J.W.* Magnetic resonance-guided focused ultrasound in neurosurgery: taking lessons from the past to inform the future. *J. Korean Med. Sci.* 2018 Oct; 33(44): e279.
- Kamel W.A., Majumdar P., Matis G. et al.* Surgical management for dystonia: efficacy of deep brain stimulation in the long term. *Neurol. Int.* 2021 Aug; 13(3): 371–386.
- Kameyama S., Shirozu H., Masuda H. et al.* MRI-guided stereotactic radiofrequency thermocoagulation for 100 hypothalamic hamartomas. *J. Neurosurg.* 2016 May; 124(5): 1503–1512.
- Kandel E.I.* Experience with the cryosurgical method in production of lesions of the extrapyramidal system. *Confin. Neurol.* 1965; 26(3-5): 306–309.
- Kandel E.I.* *Functional and stereotactic neurosurgery.* Boston, MA: Springer; 1989. 724 p.
- Karaczapryanov P., Titianova E.* High intensity focused ultrasound in neurology. *Neurosonol. Cerebr. Hemodynamics.* 2016 Sep; 12(2): 61–71.

- Kelly P. Pallidotomy in Parkinson's disease. *J. Neurosurg.* 1995 Jun; 36(6): 1154–1157.
- Khandhar S.M., Marks W.J. Epidemiology of Parkinson's disease. *Dis. Mon.* 2007 Apr; 53(4): 200–205.
- Kim M., Kim C.H., Jung H.H. et al. Treatment of major depressive disorder via magnetic resonance-guided focused ultrasound surgery. *Biol. Psychiatry.* 2018 Jan; 83(1): e17–e18.
- Kim M.J., Park S.H., Chang K.W. et al. Technical and operative factors affecting magnetic resonance imaging-guided focused ultrasound thalamotomy for essential tremor: experience from 250 treatments. *J. Neurosurg.* 2021 May; 135(6): 1780–1788.
- Kim S.J., Roh D., Jung H.H. et al. A study of novel bilateral thermal capsulotomy with focused ultrasound for treatment-refractory obsessive-compulsive disorder: 2-year follow-up. *J. Psychiatry Neurosci.* 2018 Aug; 43(5): 327–337.
- Kimber T.E., Tsai C.S., Semmler J. et al. Voluntary movements after pallidotomy in severe Parkinson's disease. *Brain.* 1999 May; 122(Pt. 5): 895–906.
- Kobus T., Vykhodtseva N., Pilatou N. et al. Safety validation of repeated blood-brain barrier disruption using focused ultrasound. *Ultrasound Med. Biol.* 2016 Feb; 42(2): 481–492.
- Kondziolka D., Ong J.G., Lee J.Y. et al. Gamma knife thalamotomy for essential tremor. *J. Neurosurg.* 2008 Jan; 108(1): 111–117.
- Krause P., Völzmann S., Ewert S. et al. Long-term effects of bilateral pallidal deep brain stimulation in dystonia: a follow-up between 8 and 16 years. *J. Neurol.* 2020; 267(6): 1622–1631.
- Krauss J.K. Deep brain stimulation for treatment of cervical dystonia. *Acta Neurochir. Suppl.* 2007; 97(Pt. 2): 201–205.
- Kreiner H.J. Stereotactic frame. Munich: GKS; 2003.
- Krishna V., Mindel J., Sammartino F. et al. A phase 1 open-label trial evaluating focused ultrasound unilateral anterior thalamotomy for focal onset epilepsy. *Epilepsia.* 2023 Apr; 64(4): 831–842.
- Krishna V., Sammartino F., Cosgrove R. et al. Predictors of outcomes after focused ultrasound thalamotomy. *Neurosurgery.* 2020 Aug; 87(2): 229–237.
- Laitinen L.V. Optimal target of pallidotomy: a controversy. In: *Pallidal surgery for the treatment of Parkinson's disease and movement disorders.* Krauss J.K., Grossman R.G., Jankovic J., editors. Philadelphia: Lippincott-Raven; 1998: 285–289.
- Laitinen L.V., Bergenheim A.T., Hariz M.I. Leksell's posteroventral pallidotomy in the treatment of Parkinson's disease. *J. Neurosurg.* 1992 Jan; 76(1): 53–61.
- Lang A.E., Lozano A.M., Montgomery E.B. et al. Posteroventral medial pallidotomy in advanced Parkinson's disease. *N. Engl. J. Med.* 1997 Oct; 337(15): 1036–1042.
- Lee W., Chung Y.A., Jung Y. et al. Simultaneous acoustic stimulation of human primary and secondary somatosensory cortices using transcranial focused ultrasound. *BMC Neurosci.* 2016 Oct; 17(1): 68.
- Legon W., Adams S., Bansal P. et al. A retrospective qualitative report of symptoms and safety from transcranial focused ultrasound for neuromodulation in humans. *Sci. Rep.* 2020 Mar; 10(1): 5573.
- Legon W., Bansal P., Tyshynsky R. et al. Transcranial focused ultrasound neuromodulation of the human primary motor cortex. *Sci. Rep.* 2018 Jul; 8(1): 10007.
- Leksell L. [Brain surgery – parkinsonism.] (In Swedish). *Lakartidningen.* 1966 Apr; 63(16): 1543–1546.
- Lhommée E., Wojtecki L., Czernecki V. et al.; EARLYSTIM study group. Behavioural outcomes of subthalamic stimulation and medical therapy versus medical therapy alone for Parkinson's disease with early motor complications (EARLYSTIM trial): secondary analysis of an open-label randomised trial. *Lancet. Neurol.* 2018 Mar; 17(1): 223–231.
- Li H., Wu Y., Pan Y. et al. Sustained relief after cessation of subthalamic stimulation for idiopathic dystonia: a 14-year observation. *Brain Stimul.* 2021 Jul-Aug; 14(4): 938–940.

- Li J.M., Li N., Wang J. et al.* [Analysis of long-term efficacy and influencing factors of subthalamic nuclear stimulation for isolated dystonia.] (In Chinese). *Zhonghua Yi Xue Za Zhi.* 2021 Feb; 101(5): 350–354.
- Liang H., Tang J., Halliwell M.* Sonoporation, drug delivery, and gene therapy. *Proc. Inst. Mech. Eng. H.* 2010; 224(2): 343–361.
- Limousin P., Foltynie T.* Long-term outcomes of deep brain stimulation in Parkinson disease. *Nat. Rev. Neurol.* 2019 Apr; 15(4): 234–242.
- Limousin P., Krack P., Pollak P. et al.* Electrical stimulation of the subthalamic nucleus in advanced Parkinson's disease. *N. Engl. J. Med.* 1998 Oct; 339(16): 1105–1111.
- Limousin P., Pollak P., Benazzouz A. et al.* Effect on parkinsonian signs and symptoms of bilateral subthalamic nucleus stimulation. *Lancet.* 1995 Jan; 345(8942): 91–95.
- Lin C.Y., Tsai C.H., Feng L.Y. et al.* Focused ultrasound-induced blood brain-barrier opening enhanced vascular permeability for GDNF delivery in Huntington's disease mouse model. *Brain Stimul.* 2019 Sep-Oct; 12(5): 1143–1150.
- Lin F., Wu D., Yu J. et al.* Comparison of efficacy of deep brain stimulation and focused ultrasound in parkinsonian tremor: a systematic review and network meta-analysis. *J. Neurol. Neurosurg. Psychiatry.* 2021; 92(4): 434–443.
- Lindstrom P.A.* Prefrontal ultrasonic irradiation – a substitute for lobotomy. *AMA Arch. Neurol. Psychiatry.* 1954 Oct; 72(4): 339–425.
- Lipsman N., Schwartz M.L., Huang Y. et al.* MR-guided focused ultrasound thalamotomy for essential tremor: a proof-of-concept study. *Lancet. Neurol.* 2013 May; 12(5): 462–468.
- Liu Y., Zhu G., Jiang Y. et al.* Comparison of short-term stimulation of the globus pallidus interna and subthalamic nucleus for treatment of primary dystonia. *World Neurosurg.* 2019 Mar; 123: e211–e217.
- Lou J.S., Jankovic J.* Essential tremor: clinical correlates in 350 patients. *Neurology.* 1991 Feb; 41(2 (Pt. 1)): 234–238.
- Louis D.N., Perry A., Wesseling P. et al.* The 2021 WHO classification of tumors of the central nervous system: a summary. *Neuro Oncol.* 2021 Aug; 23(8): 1231–1251.
- Louis E.D.* Clinical practice. Essential tremor. *N. Engl. J. Med.* 2001 Sep; 345(12): 887–891.
- Louis E.D., Benito-León J., Bermejo-Pareja F.; Neurological Disorders in Central Spain (NEDICES) Study Group.* Self-reported depression and anti-depressant medication use in essential tremor: cross-sectional and prospective analyses in a population-based study. *Eur. J. Neurol.* 2007 Oct; 14(10): 1138e46.
- Louis E.D., Thawani S.P., Andrews H.F.* Prevalence of essential tremor in a multiethnic, community-based study in northern Manhattan, New York, NY. *Neuroepidemiology.* 2009; 32(3): 208–214.
- Lynn J.G., Zwemer R.L., Chick A.J., Miller A.E.* A new method for the generation and use of focused ultrasound in experimental biology. *J. Gen. Physiol.* 1942 Nov; 26(2): 179–193.
- Lyons K.E., Pahwa R.,* editors. *Handbook of essential tremor and other tremor disorders.* Boca Raton: Taylor & Francis Group; 2005. 416 p.
- Ma X., Li T., Du L., Han T.* Research and progress of focused ultrasound in the treatment of Alzheimer's disease. *Front. Neurol.* 2023 Dec; 14: 1323386.
- Machado H.R., Hoffman H.J., Hwang P.A.* Gelastic seizures treated by resection of a hypothalamic hamartoma. *Childs Nerv. Syst.* 1991; 7: 462–465.
- Magara A., Bühler R., Moser D. et al.* First experience with MR-guided focused ultrasound in the treatment of Parkinson's disease. *J. Ther. Ultrasound.* 2014 May; 2: 11.
- Magara A.E., Gally M.N., Moser D., Jeanmonod D.* Complete resolution of chronic cluster headache following central lateral thalamotomy using incisionless MRI-guided focused ultrasound with 6 years of follow-up: illustrative case. *J. Neurosurg. Case Lessons.* 2022 Nov; 4(22): CASE22259.

*Mainprize T., Huang Y., Alkins R. et al.* Initial experience in a pilot study of blood-brain barrier opening for chemo-drug delivery to brain tumors by MR-guided focused ultrasound. *J. Ther. Ultrasound.* 2016; 4(Suppl. 1): 17–18.

*Martínez-Fernández R., Mániz-Miró J.U., Rodríguez-Rojas R. et al.* Randomized trial of focused ultrasound subthalamotomy for Parkinson's disease. *N. Engl. J. Med.* 2020 Dec; 383(26): 2501–2513.

*Martínez-Fernández R., Mahendran S., Pineda-Pardo J.A. et al.* Bilateral staged magnetic resonance-guided focused ultrasound thalamotomy for the treatment of essential tremor: a case series study. *J. Neurol. Neurosurg. Psychiatry.* 2021a Sep; 92(9): 927–931.

*Martínez-Fernández R., Matarazzo M., Mániz-Miró J.U., Obeso J.A.* The role of focused ultrasound in the management of movement disorders: insights after 5 years of experience. *Mov. Disord. Clin. Pract.* 2021b Apr; 8(5): 681–687.

*Martin E., Jeanmonod D., Morel A. et al.* High-intensity focused ultrasound for noninvasive functional neurosurgery. *Ann. Neurol.* 2009 Dec; 66(6): 858–861.

*Meng Y., Goubran M., Rabin J.S. et al.* Blood-brain barrier opening of the default mode network in Alzheimer's disease with magnetic resonance-guided focused ultrasound. *Brain.* 2023 Mar; 146(3): 865–872.

*Menon P., Kiernan M.C., Vucic S.* Cortical hyperexcitability precedes lower motor neuron dysfunction in ALS. *Clin. Neurophysiol.* 2015 Apr; 126(4): 803–809.

*Mercado-Pimentel R., Piedimonte F., Micheli F. et al.* Successful unilateral surgical approach to internal globus pallidus and ventral intermediate nucleus of the thalamus in 3 cases of myoclonus-dystonia syndrome. *Stereotact. Funct. Neurosurg.* 2021; 99(3): 250–255.

*Merola A., Singh J., Reeves K. et al.* New frontiers for deep brain stimulation: directionality, sensing technologies, remote programming, robotic stereotactic assistance, asleep procedures, and connectomics. *Front. Neurol.* 2021 Jul; 12: 694747.

*Miller T.R., Zhuo J., Eisenberg H.M. et al.* Targeting of the dentato-rubro-thalamic tract for MR-guided focused ultrasound treatment of essential tremor. *Neuroradiol. J.* 2019 Dec; 32(6): 401–407.

*Miller W.K., Becker K.N., Caras A.J. et al.* Magnetic resonance-guided focused ultrasound treatment for essential tremor shows sustained efficacy: a meta-analysis. *Neurosurg. Rev.* 2022 Feb; 45(1): 533–544.

*Mintun M.A., Lo A.C., Duggan Evans C. et al.* Donanemab in early Alzheimer's disease. *N. Engl. J. Med.* 2021 May; 384(18): 1691–1704.

*Moan C.E., Heath R.G.* Septal stimulation for the initiation of heterosexual behavior in a homosexual male. *J. Behav. Ther. Exp. Psychiatry.* 1972; 3(1): 23–30.

*Mohadjer M., Goerke H., Milios E. et al.* Long-term results of stereotaxy in the treatment of essential tremor. *Stereotact. Funct. Neurosurg.* 1990; 54-55: 125–129.

*Monteith S., Sheehan J., Medel R. et al.* Potential intracranial applications of magnetic resonance-guided focused ultrasound surgery. *J. Neurosurg.* 2013 Feb; 118(2): 215–221.

*Montgomery S.A., Åsberg M.* A new depression scale designed to be sensitive to change. *Br. J. Psychiatry.* 1979 Apr; 134(4): 382–389.

*Monti M.M., Schnakers C., Korb A.S. et al.* Non-invasive ultrasonic thalamic stimulation in disorders of consciousness after severe brain injury: a first-in-man report. *Brain Stimul.* 2016 Nov-Dec; 9(6): 940–941.

*Movement Disorder Society Task Force on Rating Scales for Parkinson's Disease.* The Unified Parkinson's Disease Rating Scale (UPDRS): status and recommendations. *Mov. Disord.* 2003 Jul; 18(7): 738–750.

*Nambu A., Chiken S., Sano H. et al.* Dynamic activity model of movement disorders: the fundamental role of the hyperdirect pathway. *Mov. Disord.* 2023 Dec; 38(12): 2145–2150.

*Narabayashi H.* [Stereotaxic apparatus, Model II.] (In Japanese). *Psychiat. Neurol. Jpn.* 1953; 54: 669.

- Narabayashi H., Okuma T.* Procaine-oil blocking of the globus pallidus for the treatment of rigidity and tremor of parkinsonism (preliminary report). *Proc. Japan Acad.* 1953; 29(3): 134–137.
- Nasreddine Z.S., Phillips N.A., Bédirian V. et al.* The Montreal Cognitive Assessment, MoCA: a brief screening tool for mild cognitive impairment. *J. Am. Geriatr. Soc.* 2005 Apr; 53(4): 695–699.
- Nichols E., Steinmetz J.D., Vollset S.E. et al.* Estimation of the global prevalence of dementia in 2019 and forecasted prevalence in 2050: an analysis for the Global Burden of Disease Study 2019. *Lancet. Public Health.* 2022 Feb; 7(2): e105–e125.
- Niranjan A., Raju S.S., Kooshkabadi A. et al.* Stereotactic radiosurgery for essential tremor: retrospective analysis of a 19-year experience. *Mov. Disord.* 2017 May; 32(5): 769–777.
- Nüssel M., Zhao Y., Knorr C. et al.* Deep brain stimulation, stereotactic radiosurgery and high-intensity focused ultrasound targeting the limbic pain matrix: a comprehensive review. *Pain Ther.* 2022 Jun; 11(2): 459–476.
- Ochiai T.* Gamma knife thalamotomy for a medically refractory tremors: longitudinal evaluation of clinical effects and MRI response patterns. *Acta Neurochir. Suppl.* 2021; 128: 127–132.
- Okun M.S., Vitek J.L.* Lesion therapy for Parkinson's disease and other movement disorders: update and controversies. *Mov. Disord.* 2004 Apr; 19(4): 375–389.
- Osterholt T., McGurrin P., Bedard P. et al.* Thalamic tremor following focused ultrasound thalamotomy for the treatment of essential tremor. *Mov. Disord. Clin. Pract.* 2020 Oct; 8(1): 139–141.
- Ostrem J.L., San Luciano M., Dodenhoff K.A. et al.* Subthalamic nucleus deep brain stimulation in isolated dystonia: a 3-year follow-up study. *Neurology.* 2017 Jan; 88(1): 25–35.
- Ozturk S., Temel Y., Aygun D., Kocabicak E.* Deep brain stimulation of the globus pallidus internus for secondary dystonia: clinical cases and systematic review of the literature regarding the effectiveness of globus pallidus internus versus subthalamic nucleus. *World Neurosurg.* 2021 Oct; 154: e495–e508.
- Öztürk G., Taşçı İ., Samancı M.Y. et al.* Is deep brain stimulation useful in Lance–Adams syndrome? *Neurol. Asia.* 2021; 26(3): 617–620.
- Pahwa R., Lyons K.E., Wilkinson S.B. et al.* Long-term evaluation of deep brain stimulation of the thalamus. *J. Neurosurg.* 2006 Apr; 104(4): 506–512.
- Palur R.S., Berk C., Schulzer M., Honey C.R.* A meta-analysis comparing the results of pallidotomy performed with microelectrode recording or macroelectrode stimulation. *J. Neurosurg.* 2002 Jun; 96(6): 1058–1062.
- Park J., Zhang Y., Vykhodtseva N. et al.* The kinetics of blood brain barrier permeability and targeted doxorubicin delivery into brain induced by focused ultrasound. *J. Control. Release.* 2012 Aug; 162(1): 134–142.
- Park Y.S., Jung N.Y., Na Y.C., Chang J.W.* Four-year follow-up results of magnetic resonance-guided focused ultrasound thalamotomy for essential tremor. *Mov. Disord.* 2019 May; 34(5): 727–734.
- Parker E., Weidman E.K., Chazen J.L. et al.* Magnetic resonance-guided focused ultrasound for ablation of mesial temporal epilepsy circuits: modeling and theoretical feasibility of a novel non-invasive approach. *J. Neurosurg.* 2019 Jun; 133(1): 63–70.
- Patzold J.* [Physics of ultrasound, and its medical applications.] (In German). *Strahlentherapie.* 1918; 76: 4.
- Perlmutter J.S., Ushe M.* Parkinson's disease – what's the FUS? *N. Engl. J. Med.* 2020 Dec; 383(26): 2582–2584.
- Peters J., Maamary J., Kyle K. et al.* Outcomes of focused ultrasound thalamotomy in tremor syndromes. *Mov. Disord.* 2024 Jan; 39(1): 173–182.
- Peto V., Jenkinson C., Fitzpatrick R.* PDQ-39: a review of the development, validation and application of a Parkinson's disease quality of life questionnaire and its associated measures. *J. Neurol.* 1998 May; 245(Suppl. 1): S10–S14.
- Pohlman R., Richter R., Parow E.* Conduction and absorption of ultrasound waves in human tissue and its therapeutic action in sciatic and plexus neuralgia. *Deutsch. Reed. Woch.* 1939; 68: 251.

- Postuma R.B., Berg D., Stern M. et al.* MDS clinical diagnostic criteria for Parkinson's disease. *Mov. Disord.* 2015 Oct; 30(12): 1591–1601.
- Racette B.A., Dowling J., Randle J., Mink J.W.* Thalamic stimulation for primary writing tremor. *J. Neurol.* 2001 May; 248(5): 380–382.
- Rajput A.H., Rajput A.* Medical treatment of essential tremor. *J. Cent. Nerv. Syst. Dis.* 2014 Apr; 6: 29–39.
- Rasche D., Rinaldi P.C., Young R.F., Tronnier V.M.* Deep brain stimulation for the treatment of various chronic pain syndromes. *Neurosurg. Focus.* 2006; 21(6): E8.
- Ravikumar V., Parker J., Hornbeck T. et al.* Cost-effectiveness of focused ultrasound, radiosurgery, and DBS for essential tremor. *Mov. Disord.* 2017 Aug; 32(8): 1165–1173.
- Reese J.P., Dams J., Winter Y. et al.* Pharmacoeconomic considerations of treating patients with advanced Parkinson's disease. *Expert Opin. Pharmacother.* 2012 May; 13(7): 939–958.
- Ren H., Wen R., Wang W. et al.* Long-term efficacy of GPi DBS for craniofacial dystonia: a retrospective report of 13 cases. *Neurosurg. Rev.* 2022 Feb; 45(1): 673–682.
- Rezai A.R., Ranjan M., Haut M.W. et al.* Focused ultrasound-mediated blood-brain barrier opening in Alzheimer's disease: long-term safety, imaging, and cognitive outcomes. *J. Neurosurg.* 2022 Nov; 139(1): 275–283.
- Ricke V., Butts Pauly K.* MR thermometry. *J. Magn. Res. Imaging.* 2008 Feb; 27(2): 376–390.
- Rooney J.A., Nyborg W.L.* Acoustic radiation pressure in a traveling plane wave. *Am. J. Physics.* 1972 Dec; 40(12): 1825–1830.
- Samuel N., Rong Ding M.Y., Sarica C. et al.* Accelerated transcranial ultrasound neuromodulation in Parkinson's disease: a pilot study. *Mov. Disord.* 2023 Dec; 38(12): 2209–2216.
- Schaltenbrand G., Wahren W.* Atlas for stereotaxy of the human brain. 2nd ed. Stuttgart: Thieme; 1977. 84 p.
- Schlesinger I., Sinai A., Zaaroor M.* MRI-guided focused ultrasound in Parkinson's disease: a review. *Parkinsons Dis.* 2017; 2017: 8124624.
- Schneider R.B., Biglan K.M.* The promise of telemedicine for chronic neurological disorders: the example of Parkinson's disease. *Lancet. Neurol.* 2017 Jul; 16(7): 541–551.
- Schuepbach W.M., Rau J., Knudsen K. et al.; EARLYSTIM Study Group.* Neurostimulation for Parkinson's disease with early motor complications. *N. Engl. J. Med.* 2013 Feb; 368(12): 610–622.
- Scott R., Gregory R., Hines N. et al.* Neuropsychological, neurological and functional outcome following pallidotomy for Parkinson's disease. A consecutive series of eight simultaneous bilateral and twelve unilateral procedures. *Brain.* 1998 Apr; 121(Pt. 4): 659–675.
- Sepúlveda Soto M.C., Fasano A.* Essential tremor: new advances. *Clin. Park. Relat. Disord.* 2020 Dec; (3): 100031.
- Sinai A., Nassar M., Eran A. et al.* Magnetic resonance-guided focused ultrasound thalamotomy for essential tremor: a 5-year single-center experience. *J. Neurosurg.* 2019 Jul 5:1–8. doi: 10.3171/2019.3.JNS19466. Online ahead of print.
- Sinai A., Nassar M., Sprecher E. et al.* Focused ultrasound thalamotomy in tremor dominant Parkinson's disease: long-term results. *J. Parkinsons Dis.* 2022; 12(1): 199–206.
- Stiburana O., Wu L.J.C., Sheffield J.K. et al.* Motor overflow and mirror dystonia. *Parkinsonism Relat. Disord.* 2009 Dec; 15(10): 758–761.
- Spiegel E.A., Wysis H.T.* Stereoccephalotomy (thalamotomy and related procedures). Part 1. Methods and stereotaxic atlas of the human brain. New York: Grune & Stratton; 1952. 176 p.
- Srinivasan R., Ben-Pazi H., Dekker M. et al.* Telemedicine for hyperkinetic movement disorders. *Tremor Other Hyperkinet. Mov. (NY).* 2020 Feb 17: 10. doi: 10.7916/tohm.v0.698.
- Steeves T.D., Day L., Dykeman J. et al.* The prevalence of primary dystonia: a systematic review and meta-analysis. *Mov. Disord.* 2012 Dec; 27(14): 1789–1796.

- Stiell I.G., Nesbitt L.P., Nichol G. et al.; OPALS Study Group. Comparison of the Cerebral Performance Category Score and the Health Utilities Index for survivors of cardiac arrest. *Ann. Emerg. Med.* 2009 Feb; 53(2): 241–248.
- Stocchi F., Tagliati M., Olanow C.W. Treatment of levodopa-induced motor complications. *Mov. Disord.* 2008; 23(Suppl. 3): S599–S612.
- Suarez-Cedeno G., Suescun J., Schiess M.C. Earlier intervention with deep brain stimulation for Parkinson's disease. *Parkinsons Dis.* 2017; 2017: 9358153.
- Sun B., Chen S., Zhan S. et al. Subthalamic nucleus stimulation for primary dystonia and tardive dystonia. *Acta Neurochir. Suppl.* 2007; 97 (Pt. 2): 207–214.
- Sun X., Guo S., Yao J. et al. Rapid inhibition of atherosclerotic plaque progression by sonodynamic therapy. *Cardiovasc. Res.* 2019 Jan; 115(1): 190–203.
- Susheela S.P., Revannasiddaiah S., Mallarajapatna G.J., Basavalingaiah A. Robotic-arm stereotactic radiosurgery as a definitive treatment for gelastic epilepsy associated with hypothalamic hamartoma. *BMJ Case Rep.* 2013 Sep; 2013: bcr2013200538.
- Talairach J., Hecaen H., David M. et al. [Recherches sur la coagulation thérapeutique des structures sous-corticales chez l'homme.] (In French). *Rev. Neurol. (Paris)*. 1949; 81: 4–24.
- Tao A., Chen G., Deng Y., Xu R. Accuracy of transcranial sonography of the substantia nigra for detection of Parkinson's disease: a systematic review and meta-analysis. *Ultrasound Med. Biol.* 2019 Mar; 45(3): 628–641.
- Tarsy D., Vitek J.L., Starr P.A., Okun M.S., editors. *Deep brain stimulation in neurological and psychiatric disorders*. Totowa: Humana Press; 2008. 617 p.
- Tasker R.R., Organ L.W., Hawrylyshyn P. Investigation of the surgical target for alleviation of involuntary movement disorders. *Appl. Neurophysiol.* 1982; 45(3): 261–274.
- Thanou M., Gedroyc W. MRI-guided focused ultrasound as a new method of drug delivery. *J. Drug Delivery.* 2013; 2013: 616197.
- Tierney T.S., Alavian K.N., Altman N. et al. Initial experience with magnetic resonance-guided focused ultrasound stereotactic surgery for central brain lesions in young adults. *J. Neurosurg.* 2022 Jan 14:1–8. doi: 10.3171/2021.10.JNS21416. Online ahead of print.
- Tisch S. Recent advances in understanding and managing dystonia. *F1000Res.* 2018 Jul; 7: F1000 Faculty Rev-1124.
- Tisch S., Kumar K.R. Pallidal deep brain stimulation for monogenic dystonia: the effect of gene on outcome. *Front. Neurol.* 2021 Jan; 11: 630391.
- Tröster A.I., Pahwa R., Fields J.A. et al. Quality of life in Essential Tremor Questionnaire (QUEST): development and initial validation. *Parkinsonism Relat. Disord.* 2005 Sep; 11(6): 367–373.
- Tsai K.W., Chen J.C., Lai H.C. et al. The distribution of skull score and skull density ratio in tremor patients for MR-guided focused ultrasound thalamotomy. *Front. Neurosci.* 2021 May; 15: 612940.
- Tung Y.S., Vlachos F., Feshitan J.A. et al. The mechanism of interaction between focused ultrasound and microbubbles in blood-brain barrier opening in mice. *J. Acoust. Soc. Am.* 2011 Nov; 130(5): 3059–3067.
- Uchida T., Nakano M., Hongo S. et al. High-intensity focused ultrasound therapy for prostate cancer. *Int. J. Urol.* 2012 Mar; 19(3): 187–201.
- Valentino F., Cosentino G., Maugeri R. et al. Is transcranial magnetic resonance imaging-guided focused ultrasound a repeatable treatment option? Case report of a retreated patient with tremor combined with parkinsonism. *Oper. Neurosurg. (Hagerstown)*. 2020 Jun; 18(6): 577–582.
- van Dyck C.H., Swanson C.J., Aisen P. et al. Lecanemab in early Alzheimer's disease. *N. Engl. J. Med.* 2023 Jan; 388(1): 9–21.
- Vidalhet M., Vercueil L., Houeto J.L. et al. Bilateral deep-brain stimulation of the globus pallidus in primary generalized dystonia. *N. Engl. J. Med.* 2005 Feb; 352(5): 459–467.



Vingerhoets F.J.G., Villemure J.G., Temperli P. et al. Subthalamic DBS replaces levodopa in Parkinson's disease: two-year follow-up. *Neurology*. 2002 Feb; 58(3): 396–401.

Vitek J.L., Bakay R.A., Freeman A. et al. Randomized trial of pallidotomy versus medical therapy for Parkinson's disease. *Ann. Neurol*. 2003 May; 53(5): 558–569.

Wang S., Zderic V., Frenkel V. Extracorporeal, low-energy focused ultrasound for noninvasive and nondestructive targeted hyperthermia. *Future Oncol*. 2010 Sep; 6(9): 1497–1511.

Ware J., Kosinski M., Keller S. SF-36 Physical and Mental Health Summary Scales. A user's manual. Boston, MA: Health Assessment Lab; 1994. 184 p.

Wårdell K., Nordin T., Vogel D. et al. Deep brain stimulation: emerging tools for simulation, data analysis, and visualization. *Front. Neurosci*. 2022 Apr; 16: 834026.

Webb H., Lubner M.G., Hinshaw J.L. Thermal ablation. *Semin. Roentgenol*. 2011 Apr; 46(2): 133–141.

Weidman E.K., Kaplitt M.G., Strybing K., Chazen J.L. Repeat magnetic resonance imaging-guided focused ultrasound thalamotomy for recurrent essential tremor: case report and review of MRI findings. *J. Neurosurg*. 2019 Jan; 132: 211–216.

Wu J., Ross J.P., Chiu J.F. Repairable sonoporation generated by microstreaming. *J. Acoustic Soc. Am*. 2002 Mar; 111(3): 1460–1464.

Wu Y., Pan Y., Huang P. et al. Subthalamic deep brain stimulation in lingual dystonia: a case series study. *Parkinsonism Relat. Disord*. 2021 Jul; 88: 114–115.

Wu Y.S., Ni L.H., Fan R.M., Yao M.Y. Meta-regression analysis of the long-term effects of pallidal and subthalamic deep brain stimulation for the treatment of isolated dystonia. *World Neurosurg*. 2019 Sep; 129: e409–e416.

Xu D.S., Chen T., Hlubek R.J. et al. Magnetic resonance imaging-guided laser interstitial thermal therapy for the treatment of hypothalamic hamartomas: a retrospective review. *Neurosurgery*. 2018 Dec; 83(6): 1183–1192.

Yacobian T.A., Dean Fang Y.H., Gerstenecker A. et al. Brain and systemic inflammation in *de novo* Parkinson's disease. *Mov. Disord*. 2023 May; 38(5): 743–754.

Yamaguchi T., Hori T., Hori H. et al. Magnetic resonance-guided focused ultrasound ablation of hypothalamic hamartoma as a disconnection surgery: a case report. *Acta Neurochir. (Wien)*. 2020 Oct; 162(10): 2513–2517.

Young R.F., Hesselgesser R.D., Ahn E. et al. Bilateral gamma Knife thalamotomy for treatment of axial tremor. *Transl. Cancer Res*. 2014 Dec; 3(6): 525–529.

Young R.F., Li F., Vermeulen S., Meier R. Gamma Knife thalamotomy for treatment of essential tremor: long-term results. *J. Neurosurg*. 2010 Jun; 112(6): 1311–1317.

Zaaroor M., Sinai A., Goldsher D. et al. Magnetic resonance-guided focused ultrasound thalamotomy for tremor: a report of 30 Parkinson's disease and essential tremor cases. *J. Neurosurg*. 2018 Jan; 128(1): 202–210.

Zhang Y., Dong H., Xu Y. et al. External ultrasound for carotid atherosclerotic plaque treatment. *J. Ultrasound Med*. 2015 Mar; 34(3): 451–459.

Zhu J., Wang S., Xu X., Qi W. Case report. Lance–Adams syndrome: a case report and retrospective review. *Int. J. Clin. Exp. Med*. 2018; 11(11): 12752–12757.

Zigmond A.S., Snaith R.P. The hospital anxiety and depression scale. *Acta Psychiatr. Scand*. 1983 Jan; 67(6): 361–370.

Zirh A., Reich S.G., Dougherty P.M., Lenz F.A. Stereotactic thalamotomy in the treatment of essential tremor of the upper extremity: reassessment including a blinded measure of outcome. *J. Neurol. Neurosurg. Psychiatry*. 1999 Jun; 66(6): 772–775.

Zubiani A. On the application of ultrasound energy to the central nervous system. *Miner. Med*. 1951; 1: 421–436.